



INTEGRAL SOLUTION OF THE TERNARY CUBIC EQUATION

$$5(x^2 + xy^2) - 9xy + x + y + 1 = 28z^3$$

R. Anbuselvi¹ S. A. Shanmugavadivu²

¹ Associate Professor of Mathematics, ADM College for women (Autonomous), Nagapattinam, Tamilnadu, India.

² Assistant Professor of Mathematics, Thiru.Vi.Ka. Govt. Arts College, Thiruvavur, Tamilnadu, India.

ABSTRACT

The Ternary cubic equation $5(x^2 + y^2) - 9xy + x + y + 1 = 28z^3$ is considered for determining its non-zero distinct integral solutions. Employing the linear transformation $x = u + v, y = u - v$, and employing the method of factorization in complex conjugates, different patterns of integral solution to the ternary cubic equation under consideration are obtained in each pattern, interesting relations among the solutions and some special polygonal numbers like pyramidal, central pyramidal numbers are exhibited.

KEY WORDS: Ternary Cubic, Integral Solutions, Polygonal Number, Pyramidal number.

1. INTRODUCTION

Diophantine equations are numerous rich because of its variety. The determination of integral solutions for Cubic (homogeneous or non-homogeneous) diophantine equations with three variables has been an interest to mathematicians. Since antiquity as can be seen from [1-3]. In this context one may refer [4-24]. In this communication, the non-homogeneous ternary cubic diophantine equation represented by $5(x^2 + y^2) - 9xy + x + y + 1 = 28z^3$ is considered for its non-zero distinct integral solution. A few interesting relations between special polygonal numbers and Pyramidal numbers are exhibited.

NOTATIONS USED

- | | | | |
|-----|-----------|---|--|
| 1. | Pr_n | - | Pronic number of rank n |
| 2. | S_n | - | Star number of rank n |
| 3. | So_n | - | Stella octangulam number of rank n |
| 4. | OH_n | - | Octangular number of rank n |
| 5. | Gno_n | - | Gnomic number of rank n |
| 6. | Obl_n | - | Oblong number of rank n |
| 7. | Pp_n | - | Pentagonal Pyramidal number of rank n |
| 8. | CP_n^6 | - | Centred hexagonal pyramidal number |
| 9. | CP_n^3 | - | Centred hexagonal pyramidal number |
| 10. | O_n | - | Octahedral number of rank n |
| 11. | $4DF_n$ | - | Four dimensional figurate number |
| 12. | $T_{m,n}$ | - | Polygonal number of rank n with size m |

2. Methods of Analysis

The Ternary Cubic Diophantine equation under consideration is

$$5(x^2 + y^2) - 9xy + x + y + 1 = 28z^3 \quad (1)$$

$$\text{Introduction of the transformations } x = u + v, y = u - v \quad (2)$$

In (1) leads to

$$(u + 1)^2 + 19v^2 = 28z^3 \quad (3)$$

Equation (3) is solved through different methods and thus, we obtain different patterns of Solutions to (1).

2.1 Method 1

$$\text{Take } z = a^2 + 19b^2 \quad (4)$$

$$\text{Write } 28 = (3 + i\sqrt{19})(3 - i\sqrt{19}) \quad (5)$$

Using (4) and (5) in (3) and employing the method of factorization, define

$$(u + 1) + i\sqrt{19}v = (3 + i\sqrt{19})(a + i\sqrt{19}b)^3 \quad (6)$$

Equating real and imaginary parts of (6) on both sides, we get

$$u + 1 = 3a^3 - 57a^2b - 171ab^2 + 361b^3$$

$$v = a^3 + 9a^2b - 57ab^2 - 57b^3$$

Substituting the values of u, v in (2), the non-zero distinct integralsolutions to (1) are given by

$$x = 4a^3 - 48a^2b - 228ab^2 + 304b^3 - 1$$

$$y = 2a^3 - 66a^2b - 114ab^2 + 418b^3 - 1$$

$$z = a^2 - 19b^2$$

PROPERTIES

- $x(a, 1) - y(a, 1) - So_n - 2t_{4,3a} \equiv 114 \pmod{113}$
- $x(a, 1) - 4CP_{6,a} + 48t_{4,a} - 303 \equiv 0 \pmod{228}$
- $y(a, 1) - x(a, 1) + 4CP_a^3 + 36t_{2,a} \equiv 114 \pmod{134}$
- $x(a, 1) + z(a, 1) - 4Cub_n - S_n + 53t_{4,a} \equiv 93 \pmod{16}$
- $z(a, a + 1) - 228FN_n^4 + 38CP_n^6 - 39t_{4,a} = 0$
- $x(a, a + 1) - 16So_n - 408Pr_a \equiv 303 \pmod{292}$
- $z(a, a + 1) - 5t_{4,2a} \equiv 19 \pmod{38}$

Note:

Re-writing (5) as

$$28 = (-3 + i\sqrt{19})(-3 - i\sqrt{19})$$

And proceeding as in method 1, the non-zero distinct integralsolutions to (1) are given by

$$x = -2a^3 - 66a^2b + 114ab^2 + 418b^3 - 1$$

$$y = -4a^3 - 48a^2b + 228ab^2 + 304b^3 - 1$$

$$z = a^2 + 19b^2$$

2.2 Method 2

Equation (3) can be written as

$$(u + 1)^2 + 19 = 28z^3 * 1$$

Write '1' as

$$1 = \frac{(9 + i\sqrt{19})(9 - i\sqrt{19})}{10^2}$$

Define

$$(u + 1 + i\sqrt{19}v) = \frac{(3 + i\sqrt{19})(9 + i\sqrt{19}b)^3 (9 + i\sqrt{19})}{10} \quad (7)$$

Equating real and imaginary parts, we have

$$u + 1 = \frac{1}{10} [8a^3 + 4332b^3 - 456ab^2 - 684a^2b] \quad (8)$$

$$v = \frac{1}{10} [12a^3 - 152b^3 - 684ab^2 + 24a^2b] \quad (9)$$

Since our aim is to find integer solution, assuming $a = 10A$ and $b = 10B$ in (4), (8) and (9) and substituting the values of u, v in (2), we obtain the distinct non-zero integral solutions to (1) as

$$\begin{aligned} x &= 10^2 [20A^3 + 4180B^3 - 1140AB^2 - 660A^2B] - 1 \\ y &= 10^2 [-4A^3 + 4484B^3 - 228AB^2 - 708A^2B] - 1 \\ z &= 10^2 [A^2 + 19B^2] \end{aligned}$$

PROPERTIES

1. $z(A, A(A+1)) - 22800FN_A^4 + 3800CP_A^6 + 39T_{4,10A} = 0$
2. $y(A, 1) + 600O_A + 70800(Ob1)_A \equiv 448399 \pmod{48200}$
3. $z(A, A^2) - 91200DF_A - 20t_{4,10A} = 0$
4. $z(A, A+1) + 2500Pr_A + T_{1002,A} \equiv 1900 \pmod{801}$
5. $x(A, 1) + 2000CP_n^6 + 13200T_{12,A} \equiv 417999 \pmod{166800}$
6. $x(A, 1) - 5y(A, 1) - 2880T_{4,10A} = 1823996$

2.3 Method 3

Instead of (5), 28 can be written as

$$28 = \frac{(6 + 2i\sqrt{19})(6 - 2i\sqrt{19})}{2^2} \quad (10)$$

Proceeding as in method 1 and performing some algebra, we obtain the distinct non-zero integral solutions to (1) as

$$\begin{aligned} x &= 2^2 [8A^3 + 608B^3 - 96A^2B - 456AB^2] - 1 \\ y &= 2^2 [4A^3 + 836B^3 - 132A^2B - 228AB^2] - 1 \\ z &= 2^2 [A^2 + 19B^2] \end{aligned}$$

PROPERTIES

1. $x(A, 1) + 32CP_A^6 + 192T_{6,A} \equiv 2431 \pmod{1632}$
2. $x(A, A+1) + 896SO_A + 1216Pr_A + 768P_A^5 \equiv 2431 \pmod{3360}$
3. $y(A, 1) - 4Az(A, 1) + 33T_{4,4A} \equiv 3343 \pmod{1216}$
4. $x(A, 1) - 2y(A, 1) - 672T_{4,A} = 9119$
5. $z(A, A(A+1)) - 912FN_A^4 + 152CP_A^6 + 39T_{4,2A} = 0$
6. $z(A, A+1) - 40T_{6,A} \equiv 76 \pmod{192}$

Note:

Re-write (7) as

$$(u + 1) + i\sqrt{19}v = (-3 + i\sqrt{19})(a + i\sqrt{19}b)^3 \left(\frac{9 + i\sqrt{19}}{10} \right)$$

and proceeding as in method 2, the non-zero distinct integral solution to (1) are given by

$$\begin{aligned} x &= 2^2 [-4A^3 + 836B^3 - 132A^2B + 228AB^2] - 1 \\ y &= 2^2 [-8A^3 + 608B^3 - 96A^2B + 456AB^2] - 1 \\ z &= 2^2 [A^2 + 19B^2] \end{aligned}$$

2.4 Method 4

Equation (7) can be written as

$$(u + 1) + i\sqrt{19}v = \frac{(6 + 2i\sqrt{19})}{2} (a + i\sqrt{19}b)^3 \frac{(9 + i\sqrt{19})}{10} \quad (11)$$

Equating real and imaginary Parts on both sides

$$u + 1 = \frac{1}{20} [16a^3 + 8664b^3 - 1368a^2b - 912ab^2] \quad (12)$$

$$v = \frac{1}{20} [24a^3 - 304b^3 + 48a^2b - 1368ab^2] \quad (13)$$

Since our aim is to find integer solutions, assuming $a=20A$, $b=20B$ in (4), (12) and (13) and substituting the values of u, v in (2), we obtain the distinct non-zero integral solutions to (1) as

$$\begin{aligned} x &= 20^2 [40A^3 + 8360B^3 - 1320A^2B - 2280AB^2] - 1 \\ y &= 20^2 [-8A^3 + 8968B^3 - 1416A^2B + 456AB^2] - 1 \\ z &= 20^2 [A^2 + 19B^2] \end{aligned}$$

PROPERTIES

1. $x(A, 1) - 40Az(A, 1) + 105600T_{12,A} \equiv 3343999 \pmod{1321600}$
2. $z(A, A+1) + 4000T_{6,A} \equiv 7600 \pmod{19200}$
3. $y(A, 1) - 1600SO_A + 1416T_{4,20A} \equiv 3587199 \pmod{180800}$
4. $x(A, 1) - y(A, 1) - 2880OH_A - 9600T_{10,A} \equiv 243200 \pmod{1075200}$
5. $x(A, A+1) - 3840000PP_A - 3600T_{4,40A} - 4560000Gno_A = 37999999$

Conclusion

In this paper, we have made an attempt to obtain a complete set of non-trivial distinct integral solutions for the non-homogeneous ternary cubic equation. To conclude, one may search for other choices of solutions to the considered cubic equation and further cubic equations with multivariables.

REFERENCES

- [1] Dickson, L.E. "History of Theory of Numbers and Diophantine Analysis", Vol.2, Dover Publications, New York 2005.
- [2] Mordell L.J., "Diophantine Equations" Academic Press, New York, 1970
- [3] RD. Carmichael, "The Theory of Numbers and Diophantine Analysis", Dover Publications, New York 1959
- [4] M.A. Gopalan, ManjuSomnath and N.Vanitha, "On ternary cubic Diophantine equation $x^2 + y^2 = z^3$ Advances in Theoretical and Applied Mathematics", Vol. 1, no.3, 227-231, 2006.
- [5] M.A. Gopalan, ManjuSomnath and N. Vanitha, "Ternary cubic Diophantine equation $x^2 - y^2 = z^3$ ", ActaCienciaIndica, Vol.XXXIIM, No. 3, 705-707, 2007.
- [6] M.A. Gopalan and R.Anbuselvi, "Integral Solutions of ternary cubic Diophantine equation $x^2 + y^2 + 4N = zxy$ ", Pure and Applied Mathematics Sciences Vol.LXVII, No. 1-2, 107-111, March 2008.
- [7] M.A. Gopalan, ManjuSomnath and N. Vanitha, "Ternary cubic Diophantine equation $22a-1(x^2+y^2)=z^3$, actaCienciaIndia, Vol.XXXIVM, No. 3, 1135-1137, 2008.
- [8] M.A. Gopalan, J.Kaligarani Integral solutions $x^3 + y^3 + 8k(x+y) = (2k+1)z^3$ Bulletin of Pure and Applied Sciences, vol 29E, No.1, 95-99,2010
- [9] M.A.Gopalan, J.Kaligarani Integral solutions $x^3 + y^3 + 8k(x+y) = (2k+1)z^3$ Bulletin of Pure and Applied Sciences, Vol. 29E, No.1, 95-99, 2010
- [10] M.A.Gopalan, S.Premalatha On the ternary cubic equation $x^3 + x^2 + y^3 - y^2 = 4(z^3 + z^2)$ Cauvery Research Journal Vol. 4, iss 1&2 87-89, July 2010-2011.
- [11] M.A.Gopalan, V.Pandichelvi, observations on the ternary cubic diophantine equation $x^3 + y^3 + x^2 - y^3 = 4(z^3 + z^2)$ Archimedes J.Math 1(1), 31-37, 2011
- [12] M.A.Gopalan, G.Srividhya Integral solutions of ternary cubic diophantine equation $x^3 + y^3 = z^2$ ActaCienciaIndica, Vol XXXVII No.4, 805-808, 2011
- [13] M.A.Gopalan, A.Vijayashankar, S.Vidhyalakshmi Integral solutions of ternary cubic equation $x^2 + y^2 - xy + 2(x+y+z) = (k^2+3)z^3$ Archimedes J.Math 1(1), 59-65, 2011
- [14] M.A.Gopalan, G.Sangeetha on the ternary cubic diophantine equation $y^2 = Dx^2 + z^3$ Archimedes J.Math 1(1), 7-14, 2011
- [15] M.A.Gopalan and B.Sivakami, "Integral Solutions of the ternary cubic Diophantine equation $4x^2 - 4xy + 6y^2 = [(k+1)2+5]w^3$ " Impact J.Sci. Tech., vol. 6, No.1, 15-22, 2012
- [16] M.A.Gopalan, S.Vidhyalakshmi and G.Sumathi, on the non-homogeneous equation with three unknowns " $x^3 + y^3 = 14z^3 + 3(x+y)$ " Discovery Science, Vol.2, No:4,37-39. Oct.2012
- [17] M.A.Gopalan, B.Sivakami Integral solutions of the ternary cubic equation $4x^2 - 4xy + 6y^2 = ((k+1)2+5)w^3$ Impact J. Sci. Tech., Vol 6 No. 1, 15-22, 2012
- [18] M.A.Gopalan, B.Sivakami on the ternary cubic diophantine equation $2xz = y^2(x+z)$ Bessel. J.Math 2(3), 171-177, 2012.

- [19] S.Vidhyalakshmi, T.R.Usha Rani and M.A.Gopalan Integral solutions of non-homogenous ternary cubit equation $ax^2+by^2=(a+b)z^3$ Diophantus J.Math, 2(1), 31-38, 2013
- [20] M.A.Gopalan, K.Geetha On the ternary cubic diophantine equation $x^2 + y^2 - xy = z^3$ Bessel J.Math., 3(2), 119-123, 2013
- [21] M.A.Gopalan, S.Vidhyalakshmi and A.Kavitha Observations on the ternary cubic equation $x^2 + y^2 + xy = 12z^3$ Antartica J.Math.10(5), 453-460, 2013
- [22] M.A.Gopalan, S.Vidhyalakshmi and K.Lakshmi Lattice points on the non homogeneous ternary cubic equation $x^3 + y^3 + z^3 (x+y+z) = 0$ Impact J. Sci. Tech., Vol 7 No.1, 21-25,2013
- [23] M.A.Gopalan, S.Vidhyalakshmi and K.Lakshmi Lattice points on the non homogeneous ternary cubic equation $x^3 + y^3 + z^3 (x+y+z) = 0$ Impact J. Sci. Tech., Vol 7 No.1, 51-55,2013
- [24] M.A.Gopalan, S.Vidhyalakshmi and S.Mallika on the ternary non homogeneous cubic equation $x^3 + y^3 - 3(x+y) = 2(3k^2-2)z^3$ Impact J. Sci. Tech., Vol 7 No.1, 41-55,2013